

From Molecular Structure to Prediction: Visual Reasoning in Organic Chemistry and Photochemistry

Juan Segurola

3Dresyns by Resyner Technologies S.L.

© 2026 Juan Segurola. All Rights Reserved.

Correspondence: info@3Dresyns.com · www.3Dresyns.com

HEB-003 · Independent Technical Article · Version v0.2 · 22 September 2026

Abstract

Organic chemistry and photochemistry can appear to the learner as catalogues of reactions, substituent effects and spectral values. A more compact understanding emerges when molecular structure is translated into electron-density distribution, polarity, conjugation, orbital-energy differences and accessible reaction pathways. This article develops that predictive chain using photoinitiator chemistry as a documented case study. The author's 1999 doctoral thesis provides experimental UV absorption, photophysical and photopolymerisation data for commercial Type I and Type II photoinitiators, including benzophenone derivatives, thioxanthenes, alpha-aminoalkylphenones and acylphosphine oxides. New figures are redrawn here from tabulated thesis data to show how structural modification can shift absorption maxima and alter spectral overlap with mercury-lamp emission lines. The purpose is not to reduce organic photochemistry to the rule that opposite charges attract or that more conjugation always produces a red shift. Rather, it is to show how a representation based on electron distribution and excited-state energetics constrains plausible behaviour and enables qualitative prediction. The central proposition is that understanding becomes predictive when a structural change can be followed through its electronic consequences to an observable response.

Keywords: photoinitiators; organic photochemistry; electron density; polarity; conjugation; chromophore; auxochrome; bathochromic shift; hypsochromic shift; Type I photoinitiator; Type II photoinitiator; benzophenone; thioxanthone; UV-visible absorption

1. Purpose and evidence boundary

This article is a chemistry-focused companion to HEB-002. HEB-002 develops the broader argument that connected understanding can make knowledge reconstructable [3]. HEB-003 asks what happens when that understanding becomes sufficiently structural to support prediction. The documentary core is the author's 1999 doctoral thesis. Numerical values and family comparisons in the data figures are redrawn from the thesis tables rather than digitised from the old plotted curves. Where the thesis itself presents a mechanistic interpretation, that interpretation is reported as such.

The article advances one central proposition:

structure → *electron distribution* → *state energies* → *absorption / reactivity* → *observable behaviour*

2. From reaction memorisation to electron distribution

A beginner can encounter organic chemistry as a long list of named reactions and arrows. A more transferable representation begins by asking where electron density is concentrated, where it is depleted, and which rearrangements are consistent with the structure. Bond polarity can be represented qualitatively by partial charges, for example $C\delta^+-O\delta^-$, without implying complete ionic charges. Nucleophilic and electrophilic behaviour then becomes less arbitrary because electron-rich regions interact with electron-deficient centres.

This representation is only a starting point. Resonance, orbital symmetry, steric accessibility, solvent, kinetics, thermodynamics and excited-state structure can overturn a simplistic electrostatic prediction. Its value is that it converts an arrow from a memorised symbol into a proposed redistribution of electron density.

3. Chromophores, substituents and spectral shifts

The thesis explicitly develops the connection between molecular structure and UV-visible absorption. It describes chromophores as groups of atoms whose energy spacing determines characteristic electronic transitions, and notes that structural changes alter both the energy and intensity of absorption. Substituent effects are classified as bathochromic or hypsochromic shifts and hyperchromic or hypochromic intensity changes. [1]

$$\Delta E = hc/\lambda$$

Accordingly, a lower-energy transition corresponds to a longer wavelength. Extension of conjugation can lower the HOMO-LUMO separation relevant to a transition and therefore move absorption to longer wavelength. The thesis also notes that a substituent can affect different transitions in different directions; a simple label attached to a substituent is therefore not a universal predictor.

Conceptual relation between conjugation, energy gap and absorption wavelength




Figure 1. Conceptual relation between conjugation, transition-energy separation and absorption wavelength. The left panel represents a larger HOMO-LUMO gap and shorter wavelength; the right panel represents a smaller gap and longer wavelength. Schematic only, not a quantum-chemical calculation.

4. Photoinitiators as a visual case of structural prediction

Photoinitiators are particularly instructive because their useful behaviour begins with absorption. The thesis states that the first step of photochemistry is photon absorption and that the nature of the excited state depends on molecular structure and molecular orbitals. It also distinguishes two basic free-radical families: Type I photoinitiators, which undergo direct photofragmentation, and Type II photoinitiators, which proceed through bimolecular processes such as electron/proton transfer with a co-initiator. [1]

For the learner, this creates a connected chain: the chromophore determines where light is absorbed; the excited-state landscape determines which primary processes are accessible; radical generation determines whether initiation can occur; and spectral overlap with the irradiation source constrains practical efficiency.

5. Benzophenone derivatives: visible consequences of structural substitution

Chapter X of the thesis compares several commercial Type II photoinitiators. Benzophenone derivatives show a strong $\pi \rightarrow \pi^*$ transition in the short-UV region and a much weaker $n \rightarrow \pi^*$ transition at longer wavelength. The thesis reports that para substitution can markedly alter the spectrum. Quantacure BMS, for example, shows a strong red-shifted absorption around 315 nm in isopropanol, interpreted as charge-transfer character introduced by the methylphenylthio substituent. Trigonal 12 also shows a red shift associated with extension of conjugation. [1]

The tabulated $\pi \rightarrow \pi^*$ absorption maxima in isopropanol provide a compact visual demonstration of this structural diversity:




Figure 2. Redrawn from Thesis Table X.1 [1]. Absorption maxima λ_{max} in isopropanol for selected Type II photoinitiators and related derivatives. Values are tabulated thesis data, not digitised curves.

The figure is not a ranking of practical photoinitiator performance. Absorption maximum alone does not determine initiation efficiency. The thesis itself emphasises that structural influences also control the nature of the lowest excited triplet state and radical-intermediate formation. [1]

6. Thioxanthenes: shifting useful absorption toward the near UV

Thioxanthenes provide a particularly clear bridge between structural modification and application. The thesis describes their value in pigmented systems as arising from longer-wavelength near-UV absorption, which can avoid part of the screening caused by coloured pigments. Substitution changes the detailed absorption profile, and the thesis reports variable red and blue solvent shifts consistent with mixed excited-state character. It also discusses differences between ITX and CPTX in relation to inductive and resonance effects of substituents. [1]

The important educational point is not that “thioxanthone means red-shifted” as a memorised label. It is that a fused chromophore plus substituent pattern changes the electronic structure, and therefore changes the energy and intensity of accessible transitions.

7. Type I near-UV photoinitiators and spectral overlap

The thesis also compares near-UV Type I photoinitiators such as Irgacure 369, Irgacure 907, Lucirin TPO and Irgacure 819 with the Type II thioxanthenes ITX and CPTX. Table XII.3 reports extinction behaviour at important medium-pressure mercury-lamp emission wavelengths. These discrete values are particularly useful for a new graph because they directly connect molecular absorption to a real irradiation source. [1]




Figure 3. Discrete values redrawn from Thesis Table XII.3 [1]. $\log \epsilon$ at major medium-pressure mercury-lamp emission wavelengths for selected near-UV photoinitiators. Markers are not joined because the table supplies discrete values rather than continuous spectra. The thesis table does not report a 404 nm value for Irgacure 907, so that point is intentionally absent.

This figure makes an engineering idea visually immediate. A photoinitiator need not merely absorb “in the UV”. Its absorption must overlap sufficiently with the spectral output actually available in the formulation and geometry of use. The thesis uses the same logic for pigmented inks: pigment absorption creates spectral windows, and photoinitiator systems can be designed to exploit those windows. [1,2]

8. The pigment spectral-window problem

Pigments compete with photoinitiators for incident radiation. The thesis defines the spectral window of a pigment as the region in which enough radiation is transmitted for the initiator to compete effectively for light. In the later optimisation work, the absorption properties of pigments and photoinitiators were used together to design photoinitiator blends with broadened UV-visible absorption. This is a concrete example of understanding becoming predictive: spectra are not merely recorded; they constrain formulation design. [1,2]

source emission $\cap$ pigment transmission $\cap$ photoinitiator absorption $\rightarrow$ usable excitation window

9. From spectral position to photochemical performance

A red shift can be useful, but it is not automatically beneficial. Longer-wavelength absorption may improve overlap with a source or penetrate a pigment window, yet practical initiation also depends on absorptivity, excited-state population, intersystem crossing, triplet energy and lifetime, bond cleavage or electron-transfer efficiency, oxygen quenching, radical reactivity, formulation viscosity and concentration. The thesis explicitly reports that photoinitiator activity is predominantly associated with chromophore absorption maxima while structural effects play a major role in the nature of the lowest excited triplet state and formation of radical intermediates. [1]

This prevents the educational model from becoming another hidden boundary. Structure-to-spectrum reasoning constrains possibilities; it does not eliminate the need for experiment.

10. Type I and Type II as different mechanistic representations

Type I and Type II classification is useful because it links excited-state structure to a different initiation route. Type I systems undergo unimolecular cleavage after excitation. Type II systems rely on a bimolecular encounter, commonly with a tertiary amine, followed by electron/proton transfer or hydrogen abstraction. The distinction gives the learner a mechanistic map that connects absorption to the generation of initiating radicals. [1]

The thesis also shows why classification should not be treated as the complete prediction. Within each family, substitution changes absorption, state character, radical yield, lifetime and sensitivity to oxygen. A family name is therefore a starting representation, not the final explanation. [1]

11. Polarity, charge transfer and solvent effects

The thesis discusses charge-transfer excited states and solvent-polarity effects. It reports that $n \rightarrow \pi^*$ transitions can shift to shorter wavelength in more polar solvents when the ground state is stabilised more strongly, whereas $\pi \rightarrow \pi^*$ transitions can shift to longer wavelength when the excited state is stabilised more strongly. This is another example in which the same observable - a spectral shift - cannot be predicted correctly from one memorised direction without identifying the transition and the states being stabilised. [1]

A deeper representation therefore asks: which state carries more charge separation, which state is stabilised by the environment, and how does that change ΔE ?

12. Understanding becomes predictive when the chain is complete

A useful conceptual chain for organic photochemistry is:

structure $\rightarrow$ electron distribution $\rightarrow$ conjugation / state character $\rightarrow \Delta E \rightarrow \lambda, \epsilon \rightarrow$ excited-state pathway $\rightarrow$ radicals $\rightarrow$ curing response

Each arrow can fail if an omitted variable dominates. Geometry can interrupt conjugation. A substituent can alter state ordering rather than simply extend a π system. A strong absorption can produce an unproductive state. A favourable photochemical pathway can be quenched by oxygen. A good initiator can be screened by a pigment. The chain is valuable precisely because it tells the scientist where to look when prediction and experiment diverge.

13. Implications for teaching organic chemistry and photochemistry

The chemistry examples support a general teaching strategy: reduce the number of isolated facts by showing the structural variables that generate them. The learner should not be asked only to remember whether a substituent gives a red shift, but to reason about conjugation, electron donation or withdrawal, charge-transfer character and the transition being observed. Likewise, a photoinitiator family should not be learned only as a list of commercial names, but as a set of related chromophores with characteristic excited-state routes.

- Ask where electron density is concentrated and depleted before asking where a curved arrow goes.
- Link substitution to resonance, conjugation and state-energy changes before memorising spectral shifts.
- Use absorption spectra and lamp lines together so that wavelength becomes an engineering constraint rather than an isolated number.
- Distinguish spectral position from photochemical efficiency.
- Use family classifications as maps, but test where members depart from the family pattern.

14. Conclusions

1. Organic reaction patterns become more reconstructable when expressed as changes in electron density rather than as isolated arrows.
2. Chromophore substitution can alter both the energy and intensity of absorption, producing bathochromic, hypsochromic, hyperchromic or hypochromic effects.
3. Extension of conjugation often lowers relevant transition energies and shifts absorption toward longer wavelength, but the direction and magnitude depend on the actual electronic states involved.
4. The 1999 thesis data provide experimental examples in benzophenone derivatives, thioxanthenes and near-UV Type I photoinitiators.
5. Absorption maximum is not a complete measure of photoinitiator performance; excited-state character, radical formation and formulation environment remain essential.
6. Practical photoinitiator selection depends on spectral overlap among the irradiation source, pigment transmission window and photoinitiator absorption.
7. A useful structural model becomes scientifically valuable when it makes testable predictions and remains open to correction by experiment.

The broader conclusion is:

When understanding is sufficiently structural, it becomes predictive.

References

- [1] Segurola, J. Photochemical Characteristics and Design of Synergistic Photoinitiator Combinations for UV Curable Inks and Coatings. PhD thesis, Manchester Metropolitan University, Department of Chemistry and Materials, in collaboration with Swale Process, Manchester, UK, 1999.
- [2] Segurola, J.; Allen, N. S.; Edge, M.; McMahon, A. Design of eutectic photoinitiator blends for UV/visible curable acrylated printing inks and coatings. *Progress in Organic Coatings* 37(1-2) (1999) 23-37. DOI: 10.1016/S0300-9440(99)00052-1.
- [3] Segurola, J. HEB-002 - The Hidden Boundary Problem: Understanding, Visualisation and Scientific Reasoning. 3Dresyns Independent Technical Articles, English author revision v0.2, 22 September 2026.

Knowledge boundary

The numerical graphs in this article are redrawn only from values tabulated in the 1999 thesis [1]. Figure 2 presents discrete λ_{max} values and Figure 3 presents discrete extinction data at specified mercury-lamp lines; neither should be interpreted as a reconstruction of a continuous spectrum between tabulated points. Mechanistic statements attributed to the thesis are reported within the scope of that work. The conceptual HOMO-LUMO figure is schematic and does not represent a quantum-chemical calculation of any specific photoinitiator.

Limitations

The photoinitiator data were generated with the instruments, solvents, irradiation sources and commercial materials of the doctoral study. The article does not assert that old commercial grades are compositionally identical to present-day products with similar trade names. Absorption behaviour alone cannot predict complete curing performance. No new spectroscopy, quantum-chemical calculation or kinetic experiment was performed for this article.