

Leavitt, Slipher, Lemaître and Hubble: the expanding universe

Juan Segurola

Core question

How did distance indicators, spectroscopy, radial-velocity measurements and relativistic cosmology combine to transform the universe from a largely static stellar system into an empirically expanding one?

Historical and evolutionary context

Henrietta Swan Leavitt's work on Cepheid variables culminated in the 1912 period-luminosity relation for stars in the Small Magellanic Cloud, providing a route to astronomical distance calibration. From 1912 onward Vesto Slipher measured large radial velocities for spiral nebulae using spectroscopy. Friedmann found dynamical cosmological solutions to Einstein's equations in 1922. In 1927 Georges Lemaître independently developed an expanding relativistic universe and related recession velocity to distance. Edwin Hubble's 1929 paper assembled galaxy distance estimates with radial velocities, mostly measured by Slipher and collaborators, to display an empirical velocity-distance relation.

What could communities know before this transition?

Astronomers could measure stellar positions, spectra, proper motions and some distances, and by the early twentieth century they could distinguish many stellar populations. The limitation was scale: distances beyond the Milky Way were difficult, spiral nebulae had uncertain status, and redshifts lacked a unified cosmological interpretation.

What changed?

Leavitt supplied a distance relation

Because the Cepheids in the Small Magellanic Cloud were approximately at a common distance, Leavitt could compare period and apparent brightness without first knowing absolute distances. Once calibrated, the relation became a rung in the cosmic distance ladder.

Slipher supplied systematic radial velocities

Spectroscopy converted line shifts into velocity information for spiral nebulae. Many of Slipher's measurements were large and redshifted. These data existed before a generally accepted expanding-universe interpretation.

Relativity connected observation to cosmic dynamics

Friedmann and Lemaître showed that Einstein's gravitational equations permitted non-static universes. Lemaître's 1927 work explicitly connected expanding models with nebular velocities. Hubble's 1929 analysis then made the velocity-distance relation especially visible observationally.

Why could this capability emerge?

The transition required photographic astronomy, spectroscopy, large telescopes, standard candles, precision clocks and relativity. It also required combining measurements from different observatories and people. No single dataset or theory was sufficient.

What later forms of knowledge did it enable?

Expansion became a foundation for physical cosmology, cosmic age estimates, Big Bang models and later measurements of cosmic acceleration. The same distance-redshift architecture underlies much of observational cosmology.

New limits and risks

Distance ladders are vulnerable to calibration bias, selection effects and population differences. Early galaxy samples were tiny by modern standards, and measured redshift does not by itself prove a unique cosmological model. Naming conventions can also distort

attribution by collapsing Leavitt, Slipher, Friedmann and Lemaître into a narrative centred only on Hubble.

What changed in the architecture of knowledge?

Cosmology became an **observationally constrained dynamical science**. Distances, spectra and gravitational theory could be linked into a quantitative history of the universe rather than treated as separate astronomical problems.

Loss, exclusion and uncertainty

Leavitt worked within the Harvard “computer” system, where women performed essential measurement labour under unequal institutional conditions. Slipher’s velocity data were heavily used in later analyses but often disappeared from simplified accounts. The 1927 Lemaître paper and its later English translation also have a complicated publication history. PIO-078 stops at this historical synthesis; later detector regimes belong to PIO-092.

Dependency map

- DIRECT ← PIO-070 Photography and spectroscopy: instruments that remember light
- DIRECT ← PIO-076 Einstein and spacetime
- INDIRECT → PIO-092 Modern detectors and the invisible universe

References

1. Leavitt, H. S. & Pickering, E. C. “Periods of 25 Variable Stars in the Small Magellanic Cloud.” *Harvard College Observatory Circular* 173 (1912).
2. Lemaître, G. “Republication of: A Homogeneous Universe of Constant Mass and Increasing Radius Accounting for the Radial Velocity of Extragalactic Nebulae.” Exact 1927-version English translation in *General Relativity and Gravitation* 45, 1635–1646 (2013). DOI: 10.1007/s10714-013-1548-3.
3. Hubble, E. “A relation between distance and radial velocity among extra-galactic nebulae.” *Proceedings of the National Academy of Sciences* 15(3), 168–173 (1929). DOI: 10.1073/pnas.15.3.168.
4. Kragh, H. *Cosmology and Controversy: The Historical Development of Two Theories of the Universe*. Princeton University Press (1996).